

FEDERACIÓN ESPELEOLÓGICA DE AMÉRICA LATINA Y DEL CARIBE
...DONDE LA ESPELEOLOGÍA ES SOLIDARIA



Boletín Informativo

Número 1

Comisión de Geoespeleología

Coordinador: Dr. Prof. Franco Urbani

Venezuela

ABRIL 1997

En este primer número de este Boletín Informativo presentamos un directorio preliminar que contiene las direcciones de personas e instituciones a los cuales se enviará este Boletín, algunas de ellas conocemos que trabajan o están interesadas en el tema de la geoespeleología, mientras que otras personas son los mismos delegados y corresponsales de la FEALC, a los cuales solicitamos nos indiquen nombres y direcciones de otras personas que puedan estar interesadas en el tema

Tabla de Contenido

Portada	1
Tabla de Contenido	2
Presentación	3
Agradecimiento.....	3
Directorio.....	4-5
Venezuelan Cave Minerals: a review	6-23
Acknowledgments	19
References.....	19-21
Tables	22-23



BOLETÍN INFORMATIVO DE LA COMISIÓN DE GEOSPELEOLOGÍA (Federación Espeleológica de América Latina y el Caribe -FEALC-) No. 1, Abril 1997

Coordinador: Dr. Franco Urbani

Apartado 47.334, Caracas 1041A, Venezuela

Fax: (58)-2-605.31.20. Email: furbani@reacciun.ve furbani@sagi.ucv.edu.ve

Índice

Presentación, agradecimiento	1
Directorio	2-3
Minerales de cuevas venezolanas, un breve resumen	4-22

PRESENTACIÓN

Durante el III Congreso Espeleológico de América Latina y el Caribe -FEALC-, efectuado del 2 al 8 de febrero de 1997 en la ciudad de Malargüe, Provincia de Mendoza, República Argentina, se crearon diversas comisiones de trabajo según las diversas disciplinas de la espeleología, una de ellas es la de **Geoespeleología** coordinada por el suscrito. Igualmente se acordó que los coordinadores empezaran con el establecimiento de metas concretas para ir desarrollando, así como ir reuniendo un grupo de interesados de los diversos países para aumentar la actividad científica en esta rama de la espeleología.

En este primer número de este Boletín Informativo presentamos un directorio preliminar que contiene las direcciones de personas e instituciones a los cuales se enviará este Boletín, algunas de ellas conocemos que trabajan o están interesadas en el tema de la geoespeleología, mientras que otras personas son los mismos delegados y corresponsales de la FEALC, a los cuales solicitamos nos indiquen nombres y direcciones de otras personas que puedan estar interesadas en el tema.

Para iniciar las actividades de la Comisión nos proponemos lo siguiente:

- Escribir a todas las organizaciones espeleológicas y afines de América Latina y el Caribe y espeleólogos en general, solicitando nos informen nombres y direcciones de otras personas e instituciones posiblemente interesadas en el tema de la geoespeleología.
- Publicar este Boletín Informativo dos veces al año, aproximadamente en abril y octubre, donde publicaremos novedades de esta disciplina, bibliografía, notas varias, actualización del directorio de personal interesado en el tema.
- Ir preparando un inventario de los minerales de cuevas en los países de América Latina y el Caribe.

El suscrito esta a la disposición de todos los interesados, para analizar gratuitamente muestras de minerales de cavidades naturales o artificiales por difracción de rayos X u otros métodos si fuese necesario (microscopía electrónica de barrido, composición química, etc.).

AGRADECIMIENTO

El Coordinador de la Comisión agradece al Director de la Escuela de Geología, Minas y Geofísica, Facultad de Ingeniería, Universidad Central de Venezuela, por las facilidades prestadas para reproducir y enviar este Boletín Informativo.

DIRECTORIO DE ENVIO DE ESTE BOLETIN

ARGENTINA

Gabriel REDONTE. Grupo Espeleológico Argentino, C.C. 232. 1403 Buenos Aires.

Alberto Carlos GARRIDO. Santa Fé # 26, (8318) Plaza Huincul, 8318 Neuquén.

Carlos BENEDETTO. Instituto Argentino Investigaciones Espeleológicas (INAE). Tomasa de San Martín 752 (Barrio Parque). (5613) Malargüe - Mendoza.

Luis Hernán CARABELLI. CERMA. C.C. 71, Suc. 2, (1402) Buenos Aires. villeneu@pinos.com

Silvia BARREDO. Nicolás Avellaneda 2130 -Olivos-. Buenos Aires 1636.

Enrique F. LIPPS. José Ingenieros 5060-5164. (1605) Carapachay, Bs. As.

BOLIVIA

Rodolfo BECERRA DE LA ROCA. Sociedad Boliviana de Espeleología. Casilla 1749. La Paz.

BRASIL

Sociedade Brasileira de Espeleología. Biblioteca. Rua Minas Gerais 221. Caixa postal 56, Monte Siao 37580-000, MG. Fax: 55-35-465.2040. Email: jal@sulminas.com.

Sociedade Excursionista e Espeleologica. Escola Federale de Minas. Caixa Postal 68. Ouro preto 35400, MG.

Grupo Bambui de Pesquisas Espeleológicas. Biblioteca. Caixa Postal 488. Belo Horizonte. MG. 30123-970.

Brandini Ribeiro Luiz Felipe. Rua dos Geraneos 207. Aguas de São Pedro - SP.

COSTA RICA

Guillermo CORTES PADILLA. (Asociación Espeleológica Costarricense). Hatillo 1, Casa no. 291, San José.

CUBA

Angel GRAÑA GONZALEZ. Ave. 9A no. 28222 entre 282 y 284. Santa Fe. Playa. La Habana. Funat@artsoft.cult.cu

Roberto GUTIERREZ DOMECH. Calle 9na., no. 8402 esq. 84. Playa. La Habana 11600.

MÉXICO

Ramón Espinasa Pereña. Sociedad Mexicana de Exploraciones Subterráneas. Ingenieros No. 29, Col. Escandón, 11800 México, DF.

José PALACIOS VARGAS. a/c Lab. de Ecología y Sistemática de Microartrópodos, Dept. Biología, Fac. Ciencias, UNAM, México 04510, D.F.

PERU

Carlos MORALES BERMÚDEZ. Centro de Exploraciones Subterráneas del Perú. Av. Brasil 1815, Lima 11.

PUERTO RICO

Abel VALE. Urb. La Cumbre, Calle E Pol # 497, Box 230, San Juan, Puerto Rico 00926-5636. Email: enlacepr@caribe.net

II

ANGUILLA

Anguilla Historical Society. P.O. Box 44.

BAHAMAS

Island Caves Research Center. P.O. Box F-931.

BELICE

Logan McNATT. P.O. Box 195, Belmopan.

BERMUDA

Robert A. POWER. P.O.Box HM 1574, Hamilton HM GX.

COLOMBIA

Eliseo AMADO. Calle 34B, no. 96-19, Int. 2, Ap. 203 (A.A. 41139), Santa Fé de Bogotá.

CURAZAO

J. de Kok. Rondeklip 77.

ECUADOR

Instituto Geofísica, Escuela Politécnica Nacional, Apartado 2759, Quito.

EL SALVADOR

Div. Centro Investigaciones Geotécnicas, Ministerio de Obras Públicas. San Salvador.

GUATEMALA

Mike SHOWEROSS.. Apartado postal 343, 03901 Antigua.

GUYANA

Commissioner. Guyana Geology & Mines Commission. P.O. Box 1028. Georgetown.

HONDURAS

Unidad de Estudios y Proyectos. Ferrocarril Nacional. San Pedro de Sula.

JAMAICA

Patrick & Patrick T. EVANS. P.O. Box 1136, Kingston.

Alan FINCHAM. 261 Sturtevant Drive. Sierra Madre. CA 91024, USA.

PARAGUAY

Federico GRESLEBIN. Casilla de Correos 1604. Asunción.

REGIONES ADMINISTRATIVAS FRANCESAS (GUAYANA FRANCESA, MARTINICA, GUADALUPE, etc.). Alain GILBERT. Dirección desconocida.

REPUBLICA DOMINICANA

Domingo ABREU. Espeleogruppo. Apartado B-100. Santo Domingo.

URUGUAY

Alejandro OLMOS FLORES. Rivera 380, Durazno.

COMITÉ ASESOR

Paolo FORTI. Università di Bologna. Dipt. Geología. Via Zamboni, 67, Bologna, ITALIA. Email: Forti@geomin.unibo.it

Adolfo ERASO R. Cátedra de Hidrogeología, Esc. Tec. Sup. Ing. Minas, c/Rios Rosas no. 21. Madrid 28003. ESPAÑA. Fax: (34)-1-336.69.77. Email: aeraso@dinge.upm.es

SOCIETA SPELEOLOGICA ITALIANA. Biblioteca. Via Zamboni, 67, Bologna, ITALIA

LIBRARY OF CONGRESS. Hispanic Acquisitions section. 101 Independence Ave., SE. Washington, DC 20540-4182 USA

FEDERATION FRANCAISE DE SPELEOLOGIE. Centre Documentation. 23 rue de Nuits. 69004 LYON. FRANCIA

UIS Com. BIBLIOGRAPHY. C/O Dr. Reno Bernasconi. Hofwilstrasse 9, Postfach 63. CH-3053 Munchenbuchsee. SUISSE - SUIZA

AMERICAN GEOGRAPHICAL SOCIETY. Collection. Milwaukee Library. University of Wisconsin. P.O. Box 399. Milwaukee, WI 53201. USA

AMERICAN GEOLOGICAL INSTITUTE. Georef Library. 4220 King Street. Alexandria, VA 22302. USA

NATIONAL SPELEOLOGICAL SOCIETY. Library. 1 Cave Avenue. Huntsville, AL 35810. USA

VENEZUELAN CAVE MINERALS: A REVIEW

Franco URBANI

Universidad Central de Venezuela, Escuela de Geología, Minas y Geofísica, Caracas 1053 & Sociedad Venezolana de Espeleología, Apartado 47334, Caracas 1041A, Venezuela.

RESUMEN

Minerales de cavidades venezolanas: una revisión

Se presenta una revisión de los minerales secundarios identificados en cavidades venezolanas, señalando su localidad, una breve descripción de las características del yacimiento y su posible origen. Al igual que en todo el mundo, en cavidades de rocas carbonáticas predominan la calcita, yeso y aragonito, mientras que en cuevas de cuarcitas está el ópalo. Se reportan los siguientes grupos de minerales: carbonatos (con 6 especies), haluros (1), nitratos (2), óxidos-hidróxidos (5), fosfatos (12), silicatos (5), sulfatos (6), arseniatos (1) y minerales orgánicos (1).

Todos los minerales mencionados se forman por reacciones químicas a temperatura y presión ambiente, pero el factor más importante para explicar su diversidad es la disponibilidad de los distintos componentes químicos involucrados, algunos de los cuales pueden provenir de la disolución o meteorización química de los minerales mayoritarios de la roca caja (ejm. carbonato, sílice), otros de constituyentes minoritarios o trazas de la misma roca caja (ejm. sulfato de la oxidación de piritita), algunos del suelo suprayacente (ejm. nitratos), mientras que el fosfato y sulfato provienen principalmente de la descomposición del guano de murciélago.

Palabras claves: minerales de cuevas, mineralogía, sulfatos, silicatos, fosfatos, carbonatos

ABSTRACT

A list of Venezuelan cave minerals and the localities where they have been found is given together with a brief description of the occurrence and its possible origin. Like in other caves from carbonate rocks throughout the world the most abundant minerals are calcite, gypsum and aragonite, while opal is the most common in quartzite caves. The following mineral groups are reported: carbonates (6 species), Halides (1), nitrates (2), oxide-hydroxides (5), phosphates (12), silicates (5), sulfates (6), arseniates (1) and organic minerals (1).

All minerals species reported form at normal environmental temperature and pressure, but the most important factor that explains their diversity is the availability of the constituent chemicals. Some components come from the main bedrock constituents (e.g. carbonate, silica), while others from minor or trace bedrock minerals (e.g. sulfate from pyrite oxidation), or from the overlying soil (e.g. nitrate), while phosphate and sulfate develop mainly from the decomposition of bat guano.

Key words: cave minerals, mineralogy, sulfates, silicates, phosphates, carbonates.

INTRODUCTION

Once cave passages are formed, secondary minerals may deposit on the internal space. Such forms are called speleothems (*spelaion* = cave, *thema* = deposit) (URBANI, 1967b), so the usual stalactites and stalagmites that strike most cave visitors are speleothems subject of study by cave mineralogists. The first mineralogical work of Venezuelan speleothems is that of William B. WHITE, J. F. HAMAN & G. L. JEFFERSON (1963) in Guácharo Cave describing several types of calcite speleothems and also being the first to identify gypsum in several types of occurrences. Since 1967 different studies have been carried out by the author and more recently in cooperation with Jacques Martini and Paolo Forti. The structure of this work follows that of the book *Cave Minerals of the World* (HILL & FORTI, 1986). A summary of the reported occurrences is presented in Table 1.

CAVE MINERALS

a. CARBONATES

Carbonates are the constituents of the most frequent speleothems in caves opened in carbonate bedrock (calcite or dolomitic limestones or marbles), they appear in many different shapes.

Aragonite CaCO₃

- *Cueva de Baruta* (Mi.11), Caracas, Miranda. URBANI (1967a, 1968).

This is a unique 200 m long “crystal cave” or geode opened in January 1956 during quarry work and very soon vandalized. Aragonite was very abundant as anthodites with transparent or white needles as long as 5 cm, some transformed to dolomite. Chemical analyses show that the aragonite crystals have higher strontium content than adjacent calcite and dolomite speleothems. It is possible that this element promoted the aragonite crystallization (see under dolomite). The bedrock is a dolomitic marble. A photograph of these speleothems is shown in URBANI (1973b, fig. 13 lower part). Chemical analysis of some samples can be seen under dolomite.

- *Cueva La Guairita 2* (Mi.16), Caracas, Miranda. URBANI (1972b, 1980).

It was found in a single anthodite flower with needles of up to 15 mm long and 1 mm in diameter growing at the very tip of a 1½ cm long and upward calcite helictite.

- *Cueva Cantera Sur de Baruta* (Mi.28), Caracas, Miranda. URBANI (1967c, 1972b).

Also as a single anthodite flower developed directly on a visible fracture on the dolomitic marble bedrock, with 1½ cm diameter in its base and 2 cm long needles.

- *Cueva Ricardo Zuloaga* (Mi.42), Miranda. URBANI *et al.* (1995b).

Included in a cauliflower shaped assemblage associated with sepiolite in which the aragonite crystals appear as well formed needles.

- *Cueva de Iglesias* (Mi.50), Caracas, Miranda.

As a white powdery and irregular 1-2 mm thick crust of a few tens of cm² on the surface of an old sediment fill. Crystals are acicular.

- *Cueva La Peonía* (La.2), Lara.

Very similar to the previous occurrence but on limestone bedrock.

- *Cueva de Tiburcio*, Yaracuy. PEREIRA Leopoldo (pers. comm., 1982).

Contains aragonite in centimeter sized anthodites.

- *Cueva sin nombre al sur de Caracas*, Miranda. HERNÁNDEZ Ramón (pers. comm., 1967).

This is a small “crystal cave” or geode discovered by R. Hernández and not revisited since then. The internal surfaces are covered with abundant aragonite anthodites.

Azurite Cu₃(CO₃)₂(OH)₂

- *Cueva de Baruta* (Mi.11), Miranda. URBANI (1967a, 1972b).

Azurite and malachite were found together in a small nest 3 cm in diameter surrounded by calcite crystals on a wall of this “crystal cave”. In this locality pyrite is found in the bedrock so it seems quite possible that chalcopyrite could be present and the necessary copper provided from its oxidation.

Calcite CaCO₃

The most common cave mineral can adopt many different shapes, such as stalactites, stalagmites, coatings and crusts, conulites, coralloids, cups, draperies, flowstone, helictites, moonmilk, cave pearls, rafts, rimstone dams (gours), cave clouds, shelfstone, cave shields and spar. Very few works have been done on calcitic speleothems.

- *Cueva del Guácharo* (Mo.1).

This is the cave with the greater variety of calcite speleothems from Venezuela, but have been little studied. Stalactites, stalagmites, draperies, flowstones, rimstone dams, helictites, cave pearls, spar and cups are common in this cave. Monocrystalline stalactites and helictites were studied by WHITE *et al.* (1963). Some orange to red colored stalactites were studied by WHITE (1981), who found that the colour is due to organic compounds. Stalactites with concentric layers of calcite and detrital clay formed by periodical flooding were reported by URBANI (1969). Small stalactites of up to 3 cm long with coralloid tips were described by URBANI (1973a).

- *Cueva Alfredo Jahn* (Mi.35), Miranda. URBANI (1973a).

Pisolites of different shapes were described, some formed around small rock fragments, pieces of roots and centipedes.

- *Cueva Quebrada Marasmita* (Mi.6), Miranda. URBANI (1973a).

Here there is a rimstone dam with rims that partially cover the pool due to accumulation of raft calcite in one side due to air currents.

- *Cueva de la Discordia 1* (Fa.34), Falcón. SVE (1974a).

This cave shows the best cave shields found in Venezuela.

- *Cueva de la Gruta* (Mi.12) and *Cueva del Refugio* (Mi.30), Miranda. URBANI (1973a).

As conulites 5 to 8 cm deep.

- *Cueva del Cerro Autana* (Am.11), Amazonas. URBANI (1976).

Calcite appears in this cave as a minor phase interlayered with opal and chalcedony.

- *Cueva de Baruta* (Mi.11), Miranda. URBANI (1967c, 1968, 1970).

As bothroidal shapes with concentric laminae of radial crystals, compenetration twinning forming "lance points", dogtooth spar, flowstone and dripstone.

- *Cueva Walter Dupouy* (Mi.2), Miranda. URBANI (1972a).

It has meter-sized rimstone dams in an active stream

- *Cueva del cañón de Sorotamia* (Zu.72), Río Socuy. Zulia. LAGARDE Joris (pers. comm.)

This locality shows the best developed cave pearls found in Venezuela. There is a nest with 20 to 30 shiny pearls almost perfectly spherical and with a mean diameter of 2-4 cm.

- *Cueva Los Encantos*, Zulia. CARREÑO Rafael (pers. comm.).

In its middle passage and for about 100 m there are large, not active, rimstone dams, they may cover an area of up to approximately 15 m² and 2.5 m deep. In other locations in the cave smaller rimstone dams are also observed. Several conulites of up to 15-20 cm deep and an upper diameter of 8 cm have also been observed in this cave.

Calcite, var Lublinita CaCO₃

- *Cueva La Milagrosa* (Mo.22), Monagas. URBANI (1977d).

This variety of calcite coats the walls and ceiling in the main passage of this cave as a soft deposit with external bothroidal shape. The individual lumps are in the scale of 5-10 cm in diameter covering areas of up to 10-20 m² while the thickness varies from a few millimeters to 20 cm. Under the optical microscope and SEM it is seen that such masses are formed by needles some of which perfectly formed, with a few tens of microns long and with an almost constant diameter of near one micron. Another type also has the external form of needles but as small crystals in an *en echelon* pattern. These speleothems originated by water condensation and/or very slow seepage from the wall fractures.

Dolomite CaMg(CO₃)₂

- *Cueva de Baruta* (Mi.11), Caracas, Miranda. URBANI (1967ac, 1968).

In this "crystal cave" dolomite speleothems are less abundant than calcite and aragonite. One of the samples had a morphology of a lathlike tip and in thin section showed concentric structures with aragonite in its center (about 5% of the total sample). Other specimens did not have aragonite. The insoluble residue of these speleothems is opal. It is believed that dolomite was formed by alteration of aragonite under the influence of seeping water with high magnesium content coming from the dolomitic marble bedrock. A photograph of these speleothems is shown in URBANI (1973b, fig. 13 upper part). This is the second cave locality worldwide in which well ordered dolomite was found. The chemical composition of some samples of this assemblage is as follows:

<i>Minerals</i>	<i>Mg (%)</i>	<i>Sr (ppm)</i>
Aragonite	1.32	507
Aragonite	0.88	470
Aragonite	0.51	400
Aragonite, dolomite	9.00	216
Dolomite, aragonite	13.0	302
Dolomite	13.1	59
Calcite	0.86	23

- *Haitón de Sabana Grande* (Fa.52), San Luis, Falcón. URBANI (1977a, 1980).

Found in a moonmilk assemblage over a clay sediment, the material also contains about 40% of micron-sized gypsum crystals. The crystal size of dolomite was below the resolution of the optical microscope and its origin is not clear.

Magnesite MgCO₃

- *Cueva Quebrada Ocumarito*, Distrito Federal. URBANI (1980).

This is a 10 m long cave developed along a fault zone in serpentinite bedrock. The mineral fills the multiple fractures of the rock producing a wedging effect that promotes the passage enlargement. The crusts thickness may vary from a few millimeters up to 6 cm thickness.

Malachite Cu₂CO₃(OH)₂

- *Cueva de Baruta* (Mi.11), Caracas, Miranda. URBANI (1967a, 1972b).

Already mentioned under azurite.

b. HALIDES

Halite NaCl

- *Cueva Alfredo Jahn* (Mi.35), Birongo, Miranda. FORTI & URBANI (1996)

Forms a sub-millimeter gray coating on the walls at the far end of the Chaguaramo Saloon. It is associated with koktaite.

c. NITRATES

Nitrammite NH₄NO₃

- *Sima Aonda Superior* (Bo.54), Auyán-tepui, Bolívar. FORTI (1994).

Forms a very small patch in a well-aerated and protected niche on the wall of this quartzite cave. The components are believed to come from the decaying organic matter of the surface.

Sveite KAl₇(NO₃)₄Cl₂(OH)₁₆.8H₂O

- *Cueva del Cerro Autana* (Am.11), Amazonas. MARTINI (1980), MARTINI & URBANI (1980, 1984), URBANI (1977b).

This is the first new mineral for science discovered in a Venezuelan cave. It occurs in a quartzite cave where it forms crusts and efflorescences on the walls in patches of about 0.5 to 1 cm thickness and up to a few square meters extent. In some cases it is clearly seen that the forming solutions seeped from the bedding planes. Under the optical microscope and SEM (URBANI, 1977b), the mineral consists of an aggregate of contorted flakes. It is possible that the nitrate, chlorine and part of the potassium, come from the organic matter from the vegetation and soil at the top of the mountain, while the aluminum and potassium come from the weathering of the trace amounts of plagioclase and micas disseminated in the rock. As the solutions are very diluted its formation requires the evaporation of large amounts of water.

d. OXIDES AND HYDROXIDES

Goethite αFeO(OH)

- *Cuevas del Cerro María Luisa* and *Cueva de Conejero*, Ciudad Piar, Bolívar. URBANI (1980).

These caves are developed below the hard lateritic iron crust that covers the surface of the hill. Goethite speleothems are present as centimeter long stalactites, small stalagmites, flowstones and coatings.

- *Sima de la Lluvia de Sarisariñama* (Bo.3), Bolívar. URBANI *et al.* (1976), URBANI (1977c).

Present as coatings and stalagmites including the largest reported goethite stalagmite in the world with about 3 m height and 1 m diameter.

- * *Sima Aonda 3*, Auyán-tepui, Bolívar. URBANI (1996a)

Crust of a few square decimeters in area and up to 4 cm thick on the wall of the weathered quartzitic bedrock.

Ice H₂O

- *Cuevas de los glaciares de Timoncitos y del pico Bompland*, Mérida. PÉREZ (1978).

In 1978 several small caves were visited in those glaciers and ice stalactites were observed. Currently, most of those glaciers have melted.

Lithiophorite (Al,Li)MnO₂(OH)₂

- *Sima Menor de Sarisariñama o Martel* (Bo.2), Bolívar. URBANI *et al.* (1976), URBANI (1977bc).

Appears in wrinkled and earthy stalactites up to 15 cm long and 20 cm in diameter at its base. It is black, clearly contrasting with the reddish colour of the quartzitic bedrock. A chemical analysis gives the following components in order of abundance: MnO 42.1%, H₂O 8.4%, Al₂O₃ 6.2%, SiO₂ 5.2%, Na₂O 2.1%, P₂O₅ 1.0%, Fe₂O₃ 0.2%, Li₂O 0.2%, with trace amounts of MgO and K₂O.

Maghemite ? α-Fe₂O₃

- *Cueva La Milagrosa* (Mo.22), Monagas. URBANI (1977d, 1980).

Appears as crusts of up to 4 mm thickness covering the limestone walls with a shiny and botryoidal shape and near bat guano deposits. A partial chemical analysis shows CaO 15.2%, Fe₂O₃ 12.4%, SiO₂ 9.8 %, P₂O₅ 5.4%, Al₂O₃ 4.4 %, with smaller amounts of Na₂O, K₂O and MgO (MnO was not determined). The sample is amorphous by X rays but the heated sample studied by P. J. Bridge (Australia) suggests it to be maghemite, possibly by alteration of goethite which is also probably present.

Mn-Fe (Amorphous oxides-hydroxides)

- *Cueva El Santuario*, Trujillo. BUZIO & FORTI (1994).

Forms black pisolites in a nest of more than a hundred of them, mainly of 3-5 mm in diameter but one reached 3 cm. The origin of such rare speleothems is related to the reducing and acid environment of the forest soil surrounding this limestone cave, which allows for the migration of Fe and Mn in the reduced state and their precipitation as chelated-metallic compounds in the aerated cave conditions.

- *Cueva Alfredo Jahn* (Mi.35), Miranda. FORTI & URBANI (1996).

In this cave two different occurrences of these compounds are found: There are millimetric-sized black pisolites having iron as the major constituent element and lesser amounts of manganese, with up to 30% of organic matter, and with an XRD peak of hydroxyl-apatite. The second occurrence is under the active stream near Salón del Chaguaramo the rock is covered with red to black 1-2 mm thick crusts. They are composed mainly of iron and minor amounts of manganese, also with 20-35% organic matter and minor amounts of manganese-bearing minerals.

- *Cueva El Samán* (Zu.30), Zulia.

Has been found in two occurrence types. The first as a 0.5-1.5 mm thick dark-brown to black and shiny coating on the limestone floor and stream clasts on a passage that during the rainy season must be under water. Under the optical microscope it is seen as multilayers. A qualitative EDX analysis shows the presence mainly of Mn, Fe with minor amounts of Al, Si and Ca. The second type is mixed with hydroxyl-apatite and will be explained under this mineral.

- * *Cueva Los Encantos*, Fundo Los Encantos, Zulia.

Same as in the first type of the previous occurrences.

- * *Sumidero Los Cantos* (Zu.70), Fundo Los Encantos, Zulia

Appears as 0.5 cm thick black and earthy coating on the limestone bedrock in a place that is under water during the rainy season.

- *Cueva La Peonía* (La.2), Lara

Same as the previous occurrence but colour is rather reddish and purple.

e. PHOSPHATES

Ardealite Ca(HPO₄)(SO₄).4H₂O

- *Cueva Ricardo Zuloaga* (Mi.42), Miranda.

Identified as a constituent of a dusty deposit of decomposed bat guano associated with whitlockite and gypsum.

Brushite $\text{CaHPO}_4 \cdot 2\text{H}_2\text{O}$

- *Cueva del Indio* (Mi.24), Miranda. FORTI *et al.* (1996).

Identified in a light orange powder and very light material produced by decomposition of bat guano, associated with gypsum. The material was collected under a calcitic crust in an area previously mined for guano. This and the samples of other minerals of the apatite group from this cave were collected near the contact of the guano material with the dolomitic bedrock.

- *Cueva de San Sebastián o la Gruta de Lourdes* (Ar.3), Aragua.

Inside a mineralized bat guano deposit as in the previous locality.

- *Cueva Ricardo Zuloaga* (Mi.42), Miranda. SILVA (1995).

Same type of occurrence as the previous locality.

Carbonate-apatite $\text{Ca}_5(\text{PO}_4, \text{CO}_3)_3$

- *Cueva del Indio* (Mi.24), Miranda. URBANI (1980), FORTI *et al.* (1996).

Also as a powdery material from the decomposition of bat guano.

Carbonate-fluor-apatite $\text{Ca}_5(\text{PO}_4, \text{CO}_3)_3\text{F}$

- *Cueva del Indio* (Mi.24), Miranda. FORTI *et al.* (1996).

Same occurrence as carbonate-apatite.

Carbonate-hydroxyl-apatite $\text{Ca}_5(\text{PO}_4, \text{CO}_3)_3(\text{OH})$

- *Cueva del Indio* (Mi.24), Miranda. PALLARÉS & URBANI (1982ab).

Same occurrence as carbonate-apatite.

- *Cueva de Los Laureles* (Zu.31), Zulia.

Dark brown and earthy 0.5 cm thick coating on the limestone bedrock. Fresh bat guano is observed nearby. Associated to smaller amounts of chlor-apatite.

- *Cueva de Pardillal* (Ar.15), Aragua.

As dark-brown crusts and rough centimetric stalactites. Associated with gypsum, hydroxyl-apatite, fluor-apatite, whitlockite and calcite. This cave is inhabited by a large bat colony.

Chlor-apatite $\text{Ca}_5(\text{PO}_4)_3\text{Cl}$

- *Cueva del Indio* (Mi.24), Miranda. FORTI *et al.* (1996).

Same as explained under carbonate-fluor-apatite.

- *Cueva de Los Laureles* (Zu.31), Zulia.

Dark brown 0.5 cm thick coating on the limestone bedrock. Fresh bat guano is observed nearby. Associated to carbonate-hydroxyl-apatite which is the main phase.

Evansite $\text{Al}_3(\text{PO}_4)(\text{OH})_6 \cdot 6\text{H}_2\text{O}$ (?) Amorphous

- *Cueva de Urutany 1* (Bo.4) and *Cueva de Urutany 2* (Bo.5), Bolívar. URBANI (1977b, 1980)

Appears as rough stalagmites (the largest is 60 cm high and 40 cm diameter in its base) and also as coatings and flowstones on the walls of this quartzite cave. Colour is yellow, yellowish brown and reddish with concentric layering and flaky appearance. Under XRD appears mainly amorphous. It is very soft and when exposed for several weeks in the laboratory environment (50-60% relative humidity) it disintegrates at touch. A chemical analysis of this material gave SiO_2 11.0%, Al_2O_3 10.7%, P_2O_5 4.9%, Na_2O 0.6%, H_2O 22.2% and minor amounts of Fe_2O_3 and K_2O . The remainder, to 100%, is organic matter. Due to the amorphous nature of the mineral its identification is not certain and probably the high silica and organic matter content may be explained by the presence of allophane (hydrous aluminum silicate) and pigotite (organic mineral). The caves develop under a dense forest having a deep organic soil and contain colonies of bats of the family *Phyllostomidae*, which may account for the presence of phosphorous and organic components while Al and Si may be provided by the quartzite weathering.

- *Cueva de Aguapira 9* (Bo.17), Aguapira, Bolívar

As a flowstone on the inclined floor of the cave with an area of about 1 m². It appears in the field as a resinous material, which when dried is reddish to brown. The surface of the flowstone is covered by many small rimstone dams of 0.5 cm height and with pools of about 1 cm². The identification is not certain and the material may be a mixture of evansite-allophane-pigotite. The cave is small and unlike the Urutany caves, no signs of bats or bat guano was seen, therefore the components needed to form these minerals may come from the decaying vegetation and the weathering of the quartzite which has traces of feldspars and micas.

Fluor-apatite Ca₅(PO₄)₃F

- *Cueva del Indio* (Mi.24), Miranda. FORTI *et al.* (1996).

Same as explained under carbonate-fluor-apatite.

- * *Sumidero Los Cantos* (Zu.70), Fundo Los Encantos, Zulia.

Appears as 0.5 cm thick light brown crust protruding 1-2 cm from the limestone bedrock wall. Found on the ceiling of the cave in a place that remains as an air-bag at moments of water flood. Appears at the would-be air-water interface above which the rock is coated with calcite, while below it is covered by clay and partially decomposed organic matter. Detrital quartz and illite-mica are mixed with the phosphate.

- *Cueva de Pardillal* (Ar.15), Aragua.

In a 2 mm thick gray to brown crust associated with calcite, gypsum, hydroxyl-apatite, carbonate-hydroxyl-apatite and whitlockite.

- *Cueva Ricardo Zuloaga* (Mi.42), Miranda.

Identified as a constituent of a dusty deposit of decomposed bat guano associated with hydroxyl-apatite and gypsum.

Hydroxyl-apatite Ca₅(PO₄)₃(OH)

- *Cueva del Guácharo* (Mo.1), Monagas. URBANI (1972b, 1980).

This was the first Venezuelan cave locality where a phosphate was identified. It appears as coatings on calcite flowstones and limestone bedrock in the small passages identified as HI-4, 5 and 6 in SVE (1968). Such passages host have small bat colonies. The crusts are black to dark brown up to 6 mm thickness but usually they are at the scale of 1-3 mm.

- *Cueva del Indio* (Mi.24), Miranda. URBANI (1972b), FORTI *et al.* (1996).

As brown to black 1-3 mm thick crusts directly on the dolomitic bedrock and other calcite speleothems. The cave hosted a large bat population. This mineral and other five phosphates reported in this cave also occur in the contact between bedrock and the guano deposit.

- *Cueva El Samán* (Zu.30), Zulia.

In one of the farthest northwestern passages of this cave, now the largest in Venezuela, there is an irregular black to very dark brown mass of up to 5 cm thick and covering about 0.5 m². The XRD pattern is that of a Mn-Fe rich amorphous mineral associated with well-crystallized hydroxy-apatite. Qualitative EDX analyses show the presence of Mn, Fe, Ca, Al, P and Si.

- *Cueva Los Laureles* (Zu.31), Zulia.

Appears as dark brown to black 1-3 mm thick crusts on the limestone bedrock. Qualitative EDX analysis shows that the external surface is rich in Mn and Fe, but inside only Ca and P were detected. Under XRD the hydroxyl-apatite pattern shows broad peaks suggesting a low crystallinity.

- *Cueva Ricardo Zuloaga* (Mi.42), Miranda. URBANI (1972bd, 1996b), SILVA (1995).

As brown to black 1-3 mm thick crusts on calcite speleothems in "Salón del Arpa" and on the dolomitic marble bedrock. Gypsum is also identifiable but rather than a chemical *in situ*, it comes from the large amount of dust raised by the visitors of this very dry and dusty cave. A detailed study of the fresh to decomposed bat guano is under way and a 3 m deep exploratory pit was excavated and sampled in which this mineral has been identified associated with whitlockite and gypsum.

- *Cuevas La Torre 1 & 2* and *Cueva Narices del Diablo*, Valle de Guanape, Anzoátegui.

As brown 1-1.5 mm thick crusts on calcite speleothems and on the limestone bedrock.

- *Sima del Peñón de Ocumare*, Ocumare del Tuy, Miranda.

Same as previous occurrence.

- *Cueva del Riíto, La Aguadita, Churuguara, Falcón.*

Same as previous occurrence but yellowish brown.

- *Cueva de Lizardo o La Cuevita (Fa.26), Falcón.*

Same as previous occurrence but associated with whitlockite.

- *Cueva de San Sebastián o la Gruta de Lourdes (Ar.3), Aragua.*

As brown to black and sometimes reddish 1-3 mm thick crusts directly on the calcite bedrock. The mineral is found in the contact between the calcite bedrock and what was the former guano deposit which was exploited as fertilizer.

- *Cueva de Pararille, Santa Cruz, Churuguara, Falcón.*

As brown to black-brown 2 mm thick crusts covering a 1 m² stalagmitic form found in the higher part of an ascending passage inhabited by a large bat colony with air temperature of 35°C. Under SEM, diatoms are visible and also many filaments (fungus and algae).

- *Cueva de Pardillal (Ar.15), Aragua.*

As brown to gray 2-8 mm crusts associated with calcite, gypsum, fluor-apatite, carbonate-hydroxyl-apatite and whitlockite.

- *Cueva de Iglesias (Mi.50), Miranda.*

As dark brown to black 1-1½ mm thick crust directly on a calcite flowstone. Found in the uppermost passage inhabited by bats.

Hydroxyl-apatite, chlorian $\text{Ca}_5(\text{PO}_4)_3(\text{OH}, \text{Cl}, \text{F})$

- *Sima del Peñón de Ocumare, Miranda.*

Dark brown coating 2 mm thick on calcite flowstone. Related to bat guano.

Leucophosphite $\text{KFe}_2(\text{PO}_4)_2(\text{OH}) \cdot 2\text{H}_2\text{O}$

- *Cueva de los Culones de Caoma (DF.10), Distrito Federal. URBANI et al. (1995a).*

This is a 30 m long cave in weathered garnet amphibolite. Leucophosphite is found as roughly 0.1 mm sized rounded nodules concentricly interlayered with opal in coralloid fingers. Under the binocular microscope the mineral shows bright yellowish to red colours.

Struvite $(\text{NH}_4)\text{MgPO}_4 \cdot 6\text{H}_2\text{O}$

- * *Cueva Ricardo Zuloaga (Mi.42), Miranda.*

Identified as a constituent of a dusty deposit of decomposed bat guano associated with hydroxyl-apatite, whitlockite and gypsum.

Whitlockite $\text{Ca}_9(\text{Mg}, \text{Fe})\text{H}(\text{PO}_4)_7$

- * *Cueva Ricardo Zuloaga (Mi.42), Miranda.*

As orange brown moist and soft coatings, 5 mm thick and covering small areas of a few square centimeters on the cave walls. Under SEM it appears as an accumulation of 5-10 µm apple-shaped forms. Another type of occurrence is inside a decomposed bat guano deposit associated with gypsum, hydroxyl-apatite, brushite, ardealite and struvite.

- *Cueva de Pardillal (Ar.15), Aragua.*

Identified in a 5 cm long stalactite with concentric layers associated with gypsum and calcite. Also in dark brown crusts with calcite, gypsum, hydroxyl-apatite, fluor-apatite and carbonate-hydroxyl-apatite.

- *Cueva de Lizardo o La Cuevita (Fa.26), Falcón.*

As 1-2 mm orange-brown crusts on flowstone with hydroxyl-apatite.

- *Cueva de los Vencejos, Peñón de las Guacas, Mi.*

Associated with gypsum in a small guano deposit formed by droppings of birds (Swifts) under their nest.

f. SILICATES

Allophane (amorphous hydrous aluminum silicate)

See comments under Evansite.

Chalcedony SiO₂

- *Cueva del Cerro Autana* (Am.11), Amazonas. URBANI (1976).

This was the first Venezuelan cave where coralloid opal speleothems were studied. Under thin section crystallized fibrous quartz (chalcedony) are seen intermixed with the opal and minor calcite layers. The speleothems are active.

Opal-A SiO₂.nH₂O

- *Cueva de Baruta* (Mi.11), Miranda. URBANI (1967c, 1972b).

This is the first opal occurrence found in a Venezuelan cave but as a very inconspicuous HCl insoluble residue of dolomite speleothems.

- *Sima de la Lluvia de Sarisariñama* (Bo.3), Bolívar. URBANI (1980).

With a wide variety of coralloid and cauliflower forms of up to several square meters of coverage. Also as metric flowstones mainly covered with coralloid surfaces. The most spectacular forms are soda-straws and stalactites as much as 35 cm long with a diameter of 2.5 cm at the upper part. This stalactitic opal is translucent and glassy in comparison with the usual darker and grayish coralloids. Under SEM all show filament structures suggesting an origin induced by biologic activity (Fig. 1).

- *Sima Mayor de Sarisariñama o Humboldt* (Bo.1), Bolívar. URBANI (1977d).

Same type of occurrence as the previous locality but with rather rare stalactitic forms.

- *Sima Menor de Sarisariñama o Martel* (Bo.2), Bolívar. URBANI (1980).

Same features as Sima de la Lluvia occurrence but here a complete sequence has been documented from soft, wet and dark bacterial mats of up to 25 cm in diameter and 1-3 cm thick on the walls where some seeping water was available, to dryer parts with some small coralloids, to very dry and older parts with well-developed coralloids, suggesting a biologic induced precipitation of the opal. The bacteria were studied by KUNICKA-GOLDFINGER (1982).

- *Cueva del Cerro Autana* (Am.11), Amazonas. URBANI (1976).

See under chalcedony.

- *Cueva El Abismo* (Bo.7), Bolívar. URBANI (1977b).

As coralloid and cauliflower forms.

- *Sima Aonda Superior* (Bo.54), Bolívar. FORTI (1994).

Mainly as coralloid forms, some of several square meters in area. Here the study of a thin section of a coralloid reveals the presence of former living beings that also suggests a biologic origin.

- *Sima Aonda* (Bo.8), Bolívar. URBANI (1996ac).

At the caves of the Aonda Platform opal coralloids are always more frequently found near the entrances in the penumbra zone with algae covering parts of the bedrock. Those coralloids are the ones that show more abundant filament structures under SEM (Fig. 2, a,b,c,d), but they also occur in samples collected deeper inside the caves (Fig. 2,e,f). One SEM image (Fig. 2f) shows a tube structure which suggests that the opal precipitated around a filament when the organism was alive. Similar structures have been recently found related to the deposition of iron, manganese and gold (GHIORSE & HIRSCH, 1979; BISCHOFF *et al.*, 1992).

- *In all other caves explored in the Precambrian quartzites of the Roraima Group: Kukenán, Chimantá, Urutany, Aguapira, Tramen, Roraima, Yuruaní, Auyán-tepui.* Bolívar and Amazonas. LAGARDE J. & C. GALÁN (pers. comm.).

As coralloid speleothems. Under SEM all samples viewed show numerous organic structures, usually filament-shaped, suggesting a biologic induced origin.

- *Most caves in sandstones of the Tertiary Mirador Formation, Páramo del Tamá and Fila de Capote, Táchira-Apure: Cueva del Loto* (Ap.1) and others. URBANI (1996c), GONZÁLEZ Freddy (pers. comm.).

As millimeter-sized rough coralloids, mainly near the entrances of the caves (Fig. 3).

- *Cueva de los Culones de Caoma* (DF.10), DF. URBANI *et al.* (1995a).

Found as 0.5-1.5 cm long coralloids developed in a weathered garnet-amphibolite. See under leucophosphate.

Palygorskite $(\text{Mg,Al})_2\text{SiO}_4\text{O}_{10}(\text{OH})\cdot 4\text{H}_2\text{O}$

- *Cueva Las Úrsulas* (Mi.47), Miranda. URBANI (1975ab).

As 1 mm thick leathery sheets and crusts on the walls and fractures. This cave developed along a joint in quartz-mica-albite-schists. The sheets are light brown but also reddish due to iron-oxide staining. They are flexible and associated with minor amounts of calcite.

Sepiolite $\text{Mg}_4\text{Si}_6\text{O}_{15}(\text{OH})_2\cdot 6\text{H}_2\text{O}$

- *Cueva Ricardo Zuloaga* (Mi.42), Miranda. URBANI *et al.* (1995a).

Appears in an elongated deposit following a fracture in the ceiling of the cave. The mineral shows a cauliflower-shaped deposit in a 1 m long section and about 3 cm wide and 1-1.5 cm thick. It is soft, humid and plastic. By XRD it appears with low-crystallinity. Qualitative EDX analysis shows the presence of Si, Ca and Mg and the sample when heated to 1200°C is transformed to diopside (Ca, Mg-pyroxene). Ranging out from the moist to drier and older parts of the occurrence sepiolite becomes more crystalline and becomes associated with increasing amounts of aragonite.

g. SULFATES

Aluminite $\text{Al}_2\text{SO}_4(\text{OH})_4\cdot 7\text{H}_2\text{O}$

- *Cueva del Indio*, Pipe, D.F.

It occurs as a 4 mm thick white and chalky efflorescence on the wall of a small tectonic cave opened on quartz-feldspar metasandstone and metaconglomerate bedrock.

Ammonium-jarosite $(\text{NH}_4)\text{Fe}_3(\text{OH})_3(\text{SO}_4)_2$

- *Cueva Alfredo Jahn* (Mi.35), Miranda. FORTI & URBANI (1996).

This mineral is present in light-blue spots on the marble wallrock of the Chaguaramo Saloon, it has millimetric thickness and mixed with hydroxyl-apatite. It is likely that the ammonium required for this mineral and for koktaite is supplied by organic matter carried by the water into the cave from the soil of surrounding forest.

Bassanite $\text{CaSO}_4\cdot \frac{1}{2}\text{H}_2\text{O}$

- *Cueva de San Sebastián o la Gruta de Lourdes* (Ar.3), Aragua.

Found on top of shale fragments from a collapse inside the middle level of this cave opened in Paleocene limestones. Looks white to slightly brown efflorescences of up to three mm thickness, soft and dusty, no traces of gypsum were found. This level of the cave is very dry and shows temperatures up to 28°C. The finding and analysis took place in 1983 but in 1996 only gypsum was found in the same spot, suggesting that bassanite formed directly in this dry environment and latter hydrated to gypsum.

Epsomite $\text{MgSO}_4\cdot 7\text{H}_2\text{O}$

- *Cueva de la cantera Sur de Baruta* (Mi.28), Miranda. URBANI (1972b, 1974).

Shows very delicate white fibrous and powdery efflorescences one cm thick covering areas of several square decimeters in very dry areas near the cave entrance. The bedrock is dolomitic marble.

- *Cueva El Ermitaño* (La.1), Lara. URBANI (1967a, 1972b, 1974).

Found in two cm thick crusts of massive epsomite with coarsely fibrous nature and transparent crystals. The surface is covered by a white powdery mineral identified as hexahydrate produced by the *in situ* dehydration of epsomite.

Gypsum $\text{CaSO}_4\cdot 2\text{H}_2\text{O}$

- *Cueva del Guácharo* (Mo.1), Monagas. WHITE *et al.* (1963), URBANI (1967a).

This was the first Venezuelan cave where this common cave mineral was identified by WHITE *et al.* (1963). This mineral occurs in a very wide variety of morphologies of white to transparent crystals forming flowers, frostworks, needle-like crystals, cotton surfaces, cockscomb stalagmites, crusts on walls and

ceilings making up sparkling passages, and extraordinary “hair” hanging more than 3 m that can be moved by air blown onto them.

- *Cueva Cruzent* (Mi.37), Miranda. URBANI (1967a).

Found in several places of the main passage as gray-colored crusts of up to one cm thick with perpendicular fibers.

- *Cueva del Indio* (Mi.24), Miranda. URBANI (1972b), FORTI *et al.* (1996).

Here two types of occurrences are found, one is as very rare layers 1-2 cm thick of almost transparent fibers, the other, more common one, is as white to yellowish powdery material associated with brushite and minerals of the apatite group. Both occur inside a decomposed bat guano deposit which was exploited for fertilizer.

- *Cueva La Peonía* (La.2), Lara. SVE (1974b).

Appears as centimetric sized flowers.

- *Cueva del Peñón del Diablo* (DF.12), Distrito Federal. URBANI (1980).

As small efflorescences on the walls near the entrance. The cave is developed in calcitic marble with visible pyrite crystals.

- *Cueva El Ermitaño* (La.1), Lara. URBANI (1972b, 1974).

As small millimeter-sized white crusts on the walls.

- *Cueva pequeña de la isla El Saco*, Mochima, Sucre.

Same as previous occurrence.

- *Cueva La Milagrosa* (Mo.22), Monagas. URBANI (1977d).

As efflorescences up to 2 mm thick at prominent high evaporative places of the bedrock. Colour may vary from transparent to reddish-colored by iron oxides. Individual crystals are about 0.5 to 0.05 mm. The cave opens in the Late Cretaceous black limestones of the Querecual Formation with high pyrite content.

- *Cueva Ricardo Zuloaga* (Mi.42), Miranda. URBANI (1972b), SILVA (1995).

As the most abundant mineral related to a guano deposit and found in light-colored powdery masses and dust. Associated with phosphates.

- *Cueva de San Sebastián o la Gruta de Lourdes* (Ar.3), Aragua.

Same as the previous locality.

- *Haitón de Sabana Grande* (Fa.52), Falcón. URBANI (1977a).

Part of a moonmilk assemblage as small crystals of 10-30 μm (40%) associated with dolomite (55%).

- *Sima Mayor de Sarisariñama or Humboldt* (Bo.1), Bolívar. URBANI (1980).

Crusts of up to 2.5 cm thick with transparent to white fibers perpendicular to the cave walls.

- *Sima Aonda Superior* (Bo.54), Bolívar. FORTI (1994).

Same as previous occurrence.

- *Gruta de los Morrocoyes*, Aragua de Maturín, Monagas. URBANI (1996d).

As crusts of up to 0.5 cm thick, white and relatively hard. Also associated in trace amounts with hexahydrate. The bedrock are subarkoses of the San Juan Formation (Late Cretaceous).

- *Cueva de Iglesias* (Mi.50), Caracas, Miranda.

As a 1-2 mm thick powdery and irregular crust on the vertical wall of a 3 m trench on a clay-silt sediment fill formed about 15 years ago (30% smectite, 30% kaolinite, 25% illite-mica, 15% quartz).

- *Cueva de Pardillal* (Ar.15), Aragua.

Appears in small stalactites and millimetric wall coatings associated with calcite, hydroxyl-apatite, carbonate-hydroxyl-apatite, fluor-apatite and whitlockite.

Hexahydrate $\text{MgSO}_4 \cdot 6\text{H}_2\text{O}$

- *Cueva El Ermitaño* (La.1), Lara. URBANI (1974).

As an *in situ* dehydration product of epsomite in this very dry cave.

- *Gruta de los Morrocoyes*, Aragua de Maturín, Monagas. URBANI (1996d).

White and soft efflorescences of 5 mm thick with fibers perpendicular to the cave wall. Since no traces of epsomite have been found, it is assumed that the mineral formed directly under a highly evaporative environment and not as a dehydration product of epsomite as in the previous locality. It is associated with

traces of gypsum. The bedrock are subarkoses of the San Juan Formation (Late Cretaceous) with traces of pyrite in some siltstone-shaly interlayers.

Koktaite $(\text{NH}_4)_2\text{Ca}(\text{SO}_4)_2 \cdot \text{H}_2\text{O}$

- *Cueva Alfredo Jahn* (Mi.35), Miranda. FORTI & URBANI (1996).

Appears mixed with hydroxyl-apatite and halite in pale-gray spots on the marble bedrock. As for the source of the required ammonium see the comment under ammonium-jarosite.

h. ARSENIATES

Mangano-berzelite $(\text{Na,Ca})_2(\text{Mg,Mn})_2(\text{AsO}_4)_3$

- * *Cueva Alfredo Jahn*, Miranda. FORTI & URBANI (1996).

This is the first report of this mineral in a cave environment and is found in red to reddish-brown 1-2 mm thick coatings on the limestone bedrock under the influence of an active stream. It is associated with amorphous Fe-Mn oxide-hydroxides. The provenance of arsenic is unknown.

i. ORGANIC MINERALS

Pigotite

See comments under Evansite.

j. RELATED FORMS

Mud and sand formations

- *Cueva del Guácharo* (Mo.1), Monagas.

"Mud and sand castle" stalagmites have been observed in the Gran Salón del Derrumbe from El Guácharo cave.

- *Cueva de Baruta* (Mi.11), Miranda. URBANI (1970).

Inside a predominantly sandy deposit that fills one of the passages there are rounded concretions forming grape-shaped accumulations. The cementing material is calcite, which includes detritic quartz, microcline, muscovite, plagioclase, opaque and clay minerals.

Clay vermiculations

- *Cueva del Pio* (Mi.22), Miranda. URBANI & PEREIRA (1970), HILL & FORTI (1986: 160), HEDGES (1993: 4).

This cave showed the best vermiculations observed in Venezuela, now completely vanished due to touching and rubbing by thousand of persons that visit the cave annually. They were of the hieroglyphic type.

- *Sima del Naranjo* (Mo.41), Monagas. LAGARDE Joris (in HEDGES, 1993:5)

Here areas of about 0.5 m² are covered by hieroglyphic vermiculations very near oil-bird (*Steatornis caripensis*) nests.

- *Cueva Ricardo Zuloaga* (Mi.42), Miranda.

Irregular millimetric to centimetric sized hieroglyphic and leopard's spots types of vermiculations covering several square meters are found on the wallrock at the lower entrance of the cave. The material is composed of clay (illite-mica), some gypsum and green algae.

Rootsicles

- *Cueva de La Guairita 2* (Mi.16), Miranda

Plant roots have penetrated the first section of this small cave attaching themselves to the subvertical wallrock; subsequently they were covered by calcite. Since the original root has decayed what remains are pipe-shaped casts attached to the wall. The external diameter varies from 1 to 3 cm while the internal hole is less than 1 cm.

Carbide related formations

- *Cueva de la Brújula* (Mi.1), Miranda

This is a cave with many calcite stalactites and flowstones but now mostly inactive and in the process of degradation. During one of the 1965-1966 visits calcium carbide was dumped on the floor of the innermost part of the cave, in 1995 on such dump were observed two irregular 1.5 cm high vertical calcite fingers 3-6 mm in diameter. Under SEM the surface appears like a smooth polygonal pavement (Fig. 4, a,b), while at a broken surface it looks like dogtooth spar crystals of micrometer size (Fig. 4,c,d).

Guano-fire materials

- *Sima Fumarola de la isla de Monos* (An.5), Anzoátegui. GALÁN & GALÁN (1983).

Combustion of bat guano started in 1977 with emission of “smoke” at the upper entrance of the cave which was mainly visible during the rainy season. A year later when the cave was visited, at a depth of 0.5 m in the former bat guano deposit the temperature was above 100°C. The residual material was identified as margarite and gypsum.

- *Cueva del Peñón de las Guacas o de los Carraos* (Mi.14), Miranda.

Here a bat deposit also burned leaving irregular slags masses of up to 15 cm³. The material is very light and porous with a metallic luster and iridescent colors. It is amorphous under XRD.

MINERALS IN ARTIFICIAL CAVITIES

Aragonite CaCO₃

- *Túnel 1 de la Quebrada Mapurite*, Distrito Federal.

Chalcanthite CuSO₄.5H₂O

- *Túnel 3 de la Quebrada Mapurite*, Distrito Federal. URBANI (1980).
- *Túnel de Aroa, Yaracuy*. URBANI (1980).
- *Túnel 1, 2 y 3 de Santa Isabel*, Guárico.
- *Túnel de Uria*, Distrito Federal.

Calcite CaCO₃

- *Túnel de Carrizal*, Miranda.
- *Túnel 1 del Viaducto 1*, Caracas-La Guaira Highway, D.F.

As white stalactites of up to 52 cm long with incipient stalagmitic growth. Some other stalactites that grow on ventilation pipes and iron reinforcements are reddish dark-brown and have about 75% calcite, 15 % quartz and 10% illite-mica, the last two minerals of detritic origin. The colour is due to a high iron content as suggested by the increase in the XRD background between 10 and 25°2θ under Cu radiation.

- *Túnel 2 de Santa Isabel*, Guárico.
- Also in many highway and old railroad tunnels.

Epsomite MgSO₄.7H₂O

- *Túnel de Uria*, Distrito Federal. URBANI (1980).
- *Túnel 3 de Santa Isabel*, Guárico.

Fe rich (amorphous oxide-hydroxides)

- *Túnel 1, 2 y 3 de Santa Isabel*, Guárico.

Preliminary mossbauer spectroscopy data allows its identification as goethite.

- *Túnel de Aroa, Yaracuy*.

Goethite αFeO(OH)

- *Túnel del Junko Country Club*, Distrito Federal. URBANI (1980).
- *Túnel 1, 2 y 3 de Santa Isabel*, Guárico.

Gypsum CaSO₄.2H₂O

- *Túnel de Uria*, Distrito Federal.

Hematite $\alpha\text{-Fe}_2\text{O}_3$

- Túnel 1 del Viaducto 1, Caracas-La Guaira Highway, D.F.

Magnetite Fe_3O_4

- Túnel 1 del Viaducto 1, Caracas-La Guaira Highway, D.F.

Malachite $\text{Cu}_2\text{CO}_3(\text{OH})_2$

- Túnel 3 de Santa Isabel, Guárico.
- Túnel de Aroa, Yaracuy.

Melanterite $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$

- Túnel de Aroa, Yaracuy.
- Túnel de Uria, Distrito Federal. URBANI (1980)

Mn rich (amorphous oxide-hydroxides)

- Túnel 2 de Santa Isabel, Guárico.

Pickeringite $\text{MgAl}_2(\text{SO}_4)_4 \cdot 22\text{H}_2\text{O}$

- Túnel de Uria, Distrito Federal.
- Túnel 3 de Santa Isabel, Guárico.

Rozenite $\text{FeSO}_4 \cdot 4\text{H}_2\text{O}$

- Túnel de Aroa, Yaracuy.

Siderotile $\text{FeSO}_4 \cdot 5\text{H}_2\text{O}$

- Túnel 3 de Santa Isabel, Guárico.
- Túnel de Aroa, Yaracuy.

ACKNOWLEDGEMENTS

To Paolo Forti (Bologna), Carol A. Hill (Albuquerque) and Jacques Martini (Pretoria) for their constant help with bibliography and advice on cave mineralogy. To Profs. Virgil Winkler and Sebastián Grande for their useful comments. To Wilmer Pérez, Carlos Galán, Joris Lagarde, Francisco Herrera, and the many others who collected minerals for study and/or accompanied the author in the cave trips. Very specially to Rafael Carreño who in the past few years has been my main supplier of the interesting but non-attractive "cave crusts". To the laboratory technicians M. D. Soto and H. Fournier.

REFERENCES

- BISCHOFF G. C. O., R. R. COENRAADS & J. LUSK. 1992. Microbial accumulation of gold: an example from Venezuela. *N. Jb. Geol. Paläont. Abn.*, 185(2): 131-159.
- BUZIO A. & P. FORTI. 1994. Las pisolitas negras de la cueva El Santuario, Santa Ana, estado Trujillo, Venezuela. *Bol. Soc. Venezolana Espeleol.*, (28):13-15.
- FORTI P. 1994. Los depósitos químicos de la Sima Aonda Superior y de otras cavidades del Auyán-tepui, Venezuela. *Bol. Soc. Venezolana Espeleol.*, (28):1-4.
- , A. ROSSI & F. URBANI. 1996. I fossati della Cueva del Indio (Caracas, Venezuela). *Atti 17th Cong. Naz. Spel.*, Castelnuovo, Garfagnana, 1994, in press.
- & F. URBANI. 1996. I nuovi minerali di grotta scoperti nella Cueva Alfredo Jahn (Venezuela). *Ibidem.*
- GALÁN C. & A. GALÁN. 1983. Notas sobre la Sima fumarola de isla de Monos, N.E. de Venezuela. *Bol. Soc. Venezolana Espeleol.*, (20):3-9.
- GHIORSE W. C. & P. HIRSCH. 1979. An untrastructural study of iron and manganese deposition associated with extracellular polymers of *Pedomicrobium*-like budding bacteria. *Arch. Microbiol.*, 123: 213-226.

- HEDGES J. 1993. A review on vermiculations. *Bol. Soc. Venezolana Espeleol.*, (27):2-6.
- HILL C. & P. FORTI. 1986. *Cave minerals of the world*. National Speleological Society, USA, 238 p.
- KUNICKA - GOLDFINGER W. 1982. Preliminary observations on the microbiology of karst caves of the Sarisariñama plateau in Venezuela. *Bol. Soc. Venezolana Espeleol.*, 10(19):133-136.
- MARTINI J. 1980. Sveite, a new mineral from Autana Cave, Territorio Federal Amazonas, Venezuela. *Transactions Geological Society South Africa*, 83:239-241.
- & F. URBANI. 1980. Sveita, $KAl_7(NO_3)_4C_{12}(OH)_{16} \cdot 8H_2O$, nuevo mineral de la cueva del cerro Autana (Am.11), Territorio Federal Amazonas, Venezuela (Abstract). *XXX Conv. Anual, Asociación Venezolana para el Avance de la Ciencia (AsoVAC)*, Mérida, *Resúmenes*, p. 21.
- & ----- 1984. Sveita, un nuevo mineral de la cueva del Cerro Autana (Am.11), Territorio Federal Amazonas, Venezuela. *Bol. Soc. Venezolana Espeleol.*, (21):13-16.
- PALLARÉS M. & F. URBANI. 1982a. Mineralogía de las espeleotemas de cuevas Venezolanas (Abstract). *Acta Científica Venezolana*, 33 (supl. 1):144. Reprinted in: *Bol. Soc. Venezolana Espeleol.*, (20):97-98.
- PÉREZ F. 1978. Cuevas de hielo en el Parque Nacional 'Sierra Nevada', estado Mérida. *Bol. Soc. Venezolana Espeleol.*, 9(17): 104-106.
- & ----- 1982b. Una ocurrencia de carbonato-hidroxil-apatito en la cueva del Indio (Mi.24), estado Miranda (Abstract). *Acta Científica Venezolana*, 33(supl. 1):143.
- SILVA M. 1995. Mineralización del Guano de Murciélago de la Cueva Ricardo Zuloaga, Peñón de las Guacas, Estado Miranda (Abstract). *XLV Conv. Anual AsoVAC, USB. Acta Cient. Venez.*, 45(supl. 1):71.
- SVE - SOCIEDAD VENEZOLANA DE ESPELEOLOGÍA. 1974a. Fa.34 - Cueva La Discordia No. 1. *Bol. Soc. Venezolana Espeleol.*, 5(1):87-89.
- SVE - SOCIEDAD VENEZOLANA DE ESPELEOLOGÍA. 1974b. La.2 - Cueva La Peonía. *Bol. Soc. Venezolana Espeleol.*, 5(1):93-96.
- URBANI F. 1967a. Venezuelan Cave Minerals. *Association William Pengelly Cave Research Centre, News* (London), (9):6-7. Reprinted in: *El Guácharo*, (5):15, 1968.
- 1967b. Las Espeleotemas. *Bol. Soc. Venezolana Espeleol.*, 1(1):23-24.
- 1967c. Notas geológicas de la cueva de Baruta, Miranda (Abstract). *XVII Convención Anual AsoVAC*, Caracas. *Resúmenes*, p. 56-57.
- 1968. Calcite, aragonite and dolomite speleothems of the Baruta Cave, Venezuela (Abstract). *Geological Society of America, Annual Meeting, New Orleans. Abstracts with Program.* p. 226-227. Reprinted in: *El Guácharo*, 1(4):31; *The New York Caver*, 1(5):87. Spanish translation in: *GEOS*, 18:74-75, 1969.
- 1969. Estalactitas con capas concéntricas de calcita y arcilla, Cueva del Guácharo, Monagas. *Bol. Soc. Venezolana Espeleol.*, 2(1):9-13.
- 1970. Concreciones en los sedimentos de la cueva de Baruta, Estado Miranda. *Bol. Soc. Venezolana Espeleol.*, 3(1):5-10. [Portuguese translation: Concreções nos sedimentos da Gruta de Baruta, Estado Miranda. *Espeleología* (Brasil), 4(5-6):16-19].
- 1972a. Notas sobre la cueva Walter Dupouy (Mi.2), Capaya, Estado Miranda. *Bol. Soc. Venezolana Espeleol.*, 3(3):169-178.
- 1972b. Lista de minerales secundarios encontrados en cuevas naturales venezolanas. *El Guácharo*, 5(1-4):17; *Circular Sociedad Venezolana Geólogos*, (55):35-36.
- 1973a. Notas preliminares sobre varios tipos de espeleotemas localizadas en cuevas Venezolanas. *El Guácharo*, 6(3-4):86-97.
- 1973b. Carsos de Venezuela. Parte 2, Calizas metamórficas de la Cordillera de la Costa. *Bol. Soc. Venezolana Espeleol.*, 4(1):15-37.
- 1974. Epsomita y hexahidrita en cuevas Venezolanas. *Bol. Soc. Venezolana Espeleol.*, 5(1):5-18.
- 1975a. Palygorskita en la cueva Las Úrsulas (Mi.47), Edo. Miranda. *Bol. Soc. Venezolana Espeleol.*, 6(11):5-12.
- 1975b. Mineralogía de espeleotemas Venezolanas (Abstract). *Symposium XXXV Aniversario Sociedad Espeleológica de Cuba*. La Habana. *Resúmenes*, p. 54-55 (español) y 144-145 (inglés). Reimpreso en *Bol. Soc. Venezolana Espeleol.*, 6(12):131-132.
- 1976. Opalo, calcedonia y calcita en la cueva del Cerro Autana (Am.11), Territorio Federal Amazonas, Venezuela. *Bol. Soc. Venezolana Espeleol.*, 7(14):129-145. (Minerals from a quartzite cave).

- 1977a. Notas sobre algunas muestras de leche de luna de cuevas de Venezuela. *Bol. Soc. Venezolana Espeleol.*, 8(16):109-115.
- 1977b. Novedades sobre estudios realizados en las formas cársicas y pseudocársicas del Escudo de Guayana. *Bol. Soc. Venezolana Espeleol.*, 8(16):175-197.
- 1977c. Nuevos comentarios sobre estudios realizados en las formas cársicas de las cuarcitas del Grupo Roraima. *Bol. Soc. Venezolana Espeleol.*, 8(15):71-77.
- 1977d. Espeleotemas de calcita (Lublinita), yeso y materiales de Guano, Cueva La Milagrosa, Monagas, Venezuela. *Bol. Soc. Venezolana Espeleol.*, 8(15):5-15
- 1980. Lista de minerales secundarios encontrados en cuevas de Venezuela. *El Guácharo*, (21):44.
- 1981. Sveita, nuevo mineral de la cueva del Cerro Autana (Am.11), Territorio Federal Amazonas. *Natura* (Soc. Cienc. Nat. La Salle, Caracas), (70-71):59-61. Reprinted in: *El Guácharo*, (22):30-31.
- 1996a. Nota mineralógica preliminar sobre la plataforma de Aonda, Auyán-tepui, Bolívar. *El Guácharo* (SVE, Caracas), (38):63.
- 1996b. El efecto del guano de murciélago en la radioactividad gamma ambiental de la cueva Ricardo Zuloaga, Miranda (Abstract). *XLVI Conv. Anual AsoVAC, Barquisimeto. Acta Cient. Venez.*, 47(supl. 1):
- 1996c. Espeleotemas de ópalo de origen biogénico en cavidades desarrolladas en rocas silíceas. Bolívar y Apure (Abstract). *XLVI Conv. Anual AsoVAC, Barquisimeto. Acta Cient. Venez.*, 47(supl. 1):
- 1996d. Notas geológicas sobre la Gruta de los Morrocayos, Aragua de Maturín, Monagas. *El Guácharo* (SVE, Caracas), (38):142-149.
- , A. MANRIQUE, A. COLOMINE, D. SOTO & S. CAMERO. 1995a. Espeleotemas de ópalo y leucofosfita en la Cueva de los Culones de Caoma, Carayaca, Distrito Federal (Abstract). *XLV Conv. Anual AsoVAC, USB. Acta Cient. Venez.*, 46(supl. 1):68.
- & J. PEREIRA. 1970. Mi.22 - Cueva del Pío. *Bol. Soc. Venezolana Espeleol.*, 3(1):24-26.
- , D. SOTO & E. DELGADO. 1995b. Una ocurrencia de sepiolita en la Cueva Ricardo Zuloaga, Peñón de las Guacas, Miranda (Abstract). *XLV Conv. Anual AsoVAC, USB. Acta Cient. Venez.*, 46(supl. 1):68-69.
- , P. ZAWIDSKY & B. KOISAR. 1976. Observaciones geológicas de la meseta de Sarisariñama, Estado Bolívar. *Boletín Informativo Asociación Venezolana Geología, Minería y Petróleo*, 19(2):77-86.
- WHITE W. B. 1981. Reflectance spectra and colour in speleothems. *National Speleological Society Bulletin*, 44:90-97.
- , J. F. HAMAN & G. L. JEFFERSON. 1963. Note on the mineralogy of Cueva del Guácharo. *Bol. Soc. Venezolana Ciencias Naturales*, 25(106):155-162.

LEYENDAS DE FIGURAS (Solicitarlas al autor)

- Fig. 1. Scanning Electron Microscope (SEM) images from the interior of an opal-A stalactite from the Sima de la Lluvia.
- Fig. 2. SEM images of opal-A coralloids from Sima Aonda near the entrance of the Alí Primera passage. Notice the residual structures of biologic origin (algae, fungus and bacteria). On image f only a pipe is seen which originally formed around and organic filament.
- Fig. 3. SEM images of opal coralloids from Cueva del Loto (Ap.1). Residual forms after bacteria and fungus are shown.
- Fig. 4. SEM images of calcite recrystallized from a 20 years old calcium carbide dump in Cueva de La Brújula. Images a and b are from the external surface, while c and d are from the interior.

Table 1 Summary of Venezuelan cave mineral occurrences (1)			#	Mor- pho- logy	Bed- rock	Environmental conditions				Provenance of chemical components										
Group	Mineral					Environmental conditions				disolution, oxidation or weathering of bedrock minerals					other sources					
						P	Aq	Guano	Evp	calcite, dolomite	Fe oxides	quartz	feldspar, micas, etc.	garnet, amphibol	serpen- tine	chalco- pyrite	pyrite	Bat guano leach.	inside	Groud w. & soil
Carbon- ates	Aragonite	CaCO ₃	8	a,e,u	C,D	X	-	-	-	Ca, CO ₃	-	-	-	-	-	-	-	-	-	
	Azurite	Cu ₃ (CO ₃) ₂ (OH) ₂	1	i	D	X	-	-	-	CO ₃	-	-	-	-	-	-	-	-	-	
	Calcite	CaCO ₃	>45 0	all	C,D	X	X	-	-	Ca, CO ₃	-	-	-	-	-	-	-	-	-	
						S	X	-	-	-	-	-	Ca	-	-	-	-	-	-	CO ₃
	Dolomite	CaMg(CO ₃) ₂	2	f,m	D,C	X	-	-	-	Ca, Mg, CO ₃	-	-	-	-	-	-	-	-	-	-
	Magnesite	MgCO ₃	1	e,r	Sr	X	-	-	-	-	-	-	-	Mg	-	-	-	-	-	-
Malachite	Cu ₂ CO ₃ (OH) ₂	1	i	D	X	-	-	-	CO ₃	-	-	-	-	-	-	-	-	-	-	
Halides	Halite	NaCl	1	e	C	X	-	-	X	-	-	-	-	-	-	-	-	-	-	
Nitrates	Nitrammite	NH ₄ NO ₃	1	e	S	X	-	-	X	-	-	-	-	-	-	-	-	-	-	NH ₄ , NO ₃
	Sveite	KAl ₇ (NO ₃) ₄ Cl ₂ (OH) ₁₆ .8H ₂ O	1	e	S	X	-	-	X	-	-	-	K, Al	-	-	-	-	-	-	Cl, NO ₃
Oxides/ Hydroxides	Goethite	alpha-FeO(OH)	4	e,s,g,f	S,Fe	X	-	-	-	-	Fe	-	-	-	-	-	-	-	-	-
	Lithiophorite	(Al,Li)MnO ₂ (OH) ₂	1	s	S	X	-	-	-	-	-	-	Al, Li	-	-	-	-	-	-	Mn?
	Maghemite ?	alpha-Fe ₂ O ₃	1	e	C	X	-	-	-	-	-	-	-	-	Fe?	-	-	-	-	Fe?
	Mn-Fe	amorphous oxides- hydroxides	6	e,p	C,D	X	X	-	-	-	-	-	-	-	Fe?	-	-	-	-	Fe, Mn
Phosphates	Ardealite	Ca(HPO ₄)(SO ₄).4H ₂ O	1	b	C,D	-	-	X	-	Ca	-	-	-	-	-	-	-	-	-	PO ₄ ,SO ₄
	Brushite	CaHPO ₄ .2H ₂ O	3	b	C,D	-	-	X	-	Ca	-	-	-	-	-	-	-	-	-	PO ₄
	Carbonate-apatite	Ca ₅ (PO ₄ , CO ₃) ₃	1	b	D	-	-	X	-	Ca, CO ₃	-	-	-	-	-	-	-	-	-	PO ₄
	Carbonate-fluor- apatite	Ca ₅ (PO ₄ ,CO ₃) ₃ F	1	b	D	-	-	X	-	Ca, CO ₃	-	-	-	-	-	-	-	-	-	F, PO ₄
	Carbonate- hydroxyl-apatite	Ca ₅ (PO ₄ ,CO ₃) ₃ (OH)	3	s,s	C	X	-	-	-	Ca, CO ₃	-	-	-	-	-	-	-	-	-	PO ₄
						b	D	-	-	X	-	-	-	-	-	-	-	-	-	PO ₄
	Chlor-apatite	Ca ₅ (PO ₄) ₃ Cl	2	e	C	X	-	-	-	Ca	-	-	-	-	-	-	-	-	-	Cl, PO ₄
						b	D	-	-	X	-	-	-	-	-	-	-	-	-	Cl, PO ₄
	Evansite	Al ₃ (PO ₄)(OH) ₆ .6H ₂ O	2	e,d	S	X	-	-	-	-	-	-	Al	-	-	-	-	-	-	PO ₄
	Fluor-apatite	Ca ₅ (PO ₄) ₃ F	2	e	C	X	-	-	-	Ca	-	-	-	-	-	-	-	-	-	PO ₄
Hydroxyl-apatite	Ca ₅ (PO ₄) ₃ (OH)	12	e,f	C,D	X	-	-	-	Ca	-	-	-	-	-	-	-	-	-	PO ₄	
					b	C,D	-	-	X	-	-	-	-	-	-	-	-	-	PO ₄	
Leucophosphite	KFe ₂ (PO ₄) ₂ (OH).2H ₂ O	1	c	Am	X	-	-	-	-	-	-	K	Fe	-	-	-	-	-	PO ₄	

	Struvite	(NH ₄)MgPO ₄ ·6H ₂ O	1	b	C,D	- - X -	Mg	- - - - -	- NH ₄ ,PO ₄	-
	Whitlockite	Ca ₉ (Mg,Fe)H(PO ₄) ₇	3	e,s	D,C	X - - -	Ca	- - - - Fe -	PO ₄	-
Silicates	Allophane	amorphous hydrous Al Si	2	e,d	S	X - - -	- - Si Al, Si	- - - -	- -	-
	Chalcedony	SiO ₂	1	s,c	S	X - - -	- - Si -	- - - -	- -	-
	Opal-A	SiO ₂ ·nH ₂ O	>30	c,s,f	S	X - - -	- - Si -	- - - -	- -	-
	Palygorskite	(Mg,Al) ₂ Si ₄ O ₁₀ (OH) ₄ ·4H ₂ O	1	r,e	Sh	X - - -	- - Si Mg, Al, Si	- - - -	- -	-
	Sepiolite	Mg ₄ Si ₆ O ₁₅ (OH) ₂ ·6H ₂ O	1	u	D	X - - -	Mg - Si	- - - -	- -	-
Sulfates	Aluminite	Al ₂ SO ₄ (OH) ₄ ·7H ₂ O	1	e	S	- - - X	- - - Al	- - SO ₄	- -	-
	Ammonium-jarosite	(NH ₄)Fe ₃ (OH) ₃ (SO ₄) ₂	1	e	D	X - - -	- - - -	- - Fe, SO ₄	- -	NH ₄
	Bassanite	CaSO ₄ ·½H ₂ O	1	e	C	X - - X	Ca - - -	- - SO ₄	- -	-
	Epsomite	MgSO ₄ ·7H ₂ O	1	e	D	X - - X	Mg - - -	- - SO ₄	- -	-
	Gypsum	CaSO ₄ ·2H ₂ O	1	e	S	X - - X	- - - -	- - SO ₄	- -	-
			10	s,l,h,i	C,D	X - - -	Ca - - -	- - SO ₄	- -	-
			3	b	C,D	- - X -	Ca - - -	- - SO ₄	- -	-
			3	e,l	S	X - - -	- - - Ca	- - SO ₄	- -	-
Hexahydrate	MgSO ₄ ·6H ₂ O	2	e	S	X - - X	Mg - - Mg	- - SO ₄	- -	-	
Koktaite	(NH ₄) ₂ Ca(SO ₄) ₂ ·H ₂ O	1	e	D	X - - -	Ca - - -	- - SO ₄	- -	NH ₄	
Arsenianterials	Manganoberzelite	(Na,Ca) ₂ (Mg,Mn) ₂ (AsO ₄) ₃	1	e	D	- X - -	Ca, Mg - - Na	- - - -	- -	Mn, As?
Organic	Pigotite	C, H, O	2	e,d	S	X - - -	- - - -	- - - -	- -	X

(1) **Abbreviations:** # = number of known occurrences of each mineral up to December 1996. **Morphology:** a= anthodite, b= crystals and powder inside bat guano, c= coralloid, d= rimstone dam, e= efflorescence, crust, coating, f= flowstone g=stalagmite, h=hair, i= isolated crystals, l= flower, m=moonmilk, p= pisolite, r= fracture fill, s= stalactite. **Bedrock types:** C= calcitic limestone or marble. D= dolomitic marble. S= sandstone or quartzite. Sh= quartz-mica-feldspar schist. Am= garnet amphibolite. Sr= serpentinite. Fe= iron laterite or iron quartzite. **Environmental conditions:** P= subaerial precipitation from dripping, flowing or seeping water. Aq= subaqueous precipitation. Guano= crystalization inside a bat guano deposit. Evp= Strong evaporative environment. **Bat guano:** Leach.= components leached from the decomposition of bat drippings or guano. Inside= components originated and staying *in situ* inside a bat guano deposit. **Ground w. & soil=** input of chemical components into the cave by water (slowly seeping, dripping, streams, etc.) and originated either from the above soil or from farther away in the drainage basin.